

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 7/64, C11B 1/10, C12N 1/06, 1/00	A1	(11) International Publication Number: WO 97/04121 (43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/GB96/01703 (22) International Filing Date: 16 July 1996 (16.07.96) (30) Priority Data: 9514649.4 18 July 1995 (18.07.95) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): LIDDELL, John, Macdonald [GB/GB]; 678 Yarm Road, Eaglescliffe TS16 0DP (GB). (74) Agents: LOCKE, Timothy, John; Zeneca LifeScience Molecules, Biomolecules, Intellectual Property Group, Belasis Avenue, P.O. Box 2, Billingham, Cleveland TS23 1YN (GB) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: EXTRACTION OF TRIGLYCERIDES FROM MICROORGANISMS (57) Abstract Oil is extracted from oil containing microorganisms by disintegrating the microorganisms and contacting them in the presence of a water content of at least 70 % by weight of that originally present in the cellular material with a water immiscible solvent for the oil, separating the solvent from the microorganisms and recovering the oil from the solvent.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

EXTRACTION OF TRIGLYCERIDES FROM MICROORGANISMS

This invention relates to an extraction process.

Vegetable oils can be extracted from dried plant seeds using pressure. This in general leaves vegetable oil in the crushed plant tissue and this may be extracted using a water-immiscible organic solvent, for example hexane.

It is possible to extract microbial oils from whole microbes by hexane extraction. The microbial material must be dried usually to a water content of less than 5% (w/w) before the oil can be effectively extracted by hexane. Not only does this involve a considerable consumption of energy because, in general, microbial cells contain at least 70% for example 80 to 95% (w/w) water based on their non-oil content, but we have also found that sensitive oils for example oxygen or heat sensitive oils may undergo chemical reactions in the drying process.

In alternative methods of extracting oils from microorganisms, water-miscible solvents, for example isopropanol, may be used. Such extractions are in general less selective than those with non-polar solvents and a wide range of other cellular components may be extracted in addition to the oils, for example phospholipids and cell wall components. Evaporation of the water-miscible solvent leaves a residue which must therefore be purified further.

By "oils" is meant materials which are liquid at the extraction temperature and which are sparingly soluble in water.

In general oils recovered from microorganisms are

triglycerides.

This invention comprises a process in which an oil is extracted from oil containing microorganisms which comprises disintegrating the microorganisms and
5 contacting them in the presence of a water content of at least 70% by weight of that originally present in the cellular material and preferably in the presence of substantially all of the original water content of the microorganisms with a water immiscible solvent for the
10 oil, separating the solvent from the microorganisms and recovering the oil from the solvent.

The invention also comprises a process for extracting a triglyceride from microbial matter containing it in which the triglyceride is released from
15 the cells by pressure homogenisation and the released triglyceride is then extracted by contacting the microbial matter with a water-immiscible solvent for example an alkane suitably having 4 to 12 and preferably 5 to 8 carbon atoms, for example cyclohexane or
20 preferably hexane. The invention avoids the requirement for dehydrating the microbial matter whilst permitting a good selectivity of extraction into the water-immiscible solvent to be secured.

It is preferred that the microorganism should be
25 disintegrated in the presence of an aqueous culture medium in which it has been cultured, as this avoids the need for separating the microbial matter from the culture medium prior to the process. Typically the organism will constitute 5% to 20% by volume of the culture medium. The
30 same solvent may be re-used two or more times for extracting successive quantities of disintegrated

organism containing triglycerides, thus increasing the triglyceride content of the solvent. This leads to economies in recovering the triglyceride from the solvent by evaporation. Such processes may, if desired, be
5 conducted by countercurrent extraction. Suitably, this may be carried out by using a solvent which is less dense than the phase containing the disintegrated microorganism and feeding it at a lower level than a phase which contains the disintegrated microorganism to a vessel in
10 which extraction occurs. The vessel may be provided with baffles, restricting vertical flow, which define contact zones between the baffles and in which means is provided for stirring material in the contact zones.

If desired, separated solvent containing the
15 triglyceride may be used together with fresh solvent to extract previously unextracted microorganisms. The solvent may be separated by centrifuging. The solvent may suitably be a hydrocarbon, for example an alkane which suitably has 4 to 12 and preferably 6 to 10 carbon
20 atoms and is suitably hexane.

The organisms may be disintegrated by enzyme cell disruption using cell wall lytic enzymes, mechanical methods such as bead milling, colloid milling, disruption by pressure release, impinging jets, ultrasonication and
25 preferably using high shear mixing and/or high pressure homogenisation. The disruption should be sufficient to disrupt the cell wall, thus enhancing the access of the solvent to the triglycerides contained within the cell. Any form of pressure homogeniser may be employed for this
30 purpose.

In general in solvent extraction processes, small

quantities of solvent are used compared with the material which is to be extracted. This has the advantage of achieving a high concentration of the desired solute in the solvent phase but when microbial materials are
5 extracted in this way there is a tendency for organic matter present to act as an emulsifying agent, leading to the formation of a solvent-in-water emulsion of high stability. In order to reduce the stability of such emulsions, it may be necessary to add demulsifying agents
10 thus enabling the emulsion to be separated suitably using high gravity conditions. However, the demulsifying agent may contaminate the product which it is desired to separate and involves additional cost. Contamination may be particularly undesirable if the demulsifier is toxic.

15 We have found that by operating in the presence of high concentrations of solvent sufficient to form a water-in-solvent dispersion, the dispersion is less stable and may be more readily separated in the absence of a demulsifying agent, for example by gravity settling
20 or use of a centrifuge. It is desirable in this case to increase the triglyceride content of the solvent by extracting successive quantities of microbial material with it.

This invention therefore comprises a process as
25 aforesaid in which the disintegrated microorganisms are contacted with a continuous phase of the solvent. It is desirable that the ratio of solvent phase to other matter present during the extraction step should be at least 1 to 1, preferably at least 1.5 to 1 and more preferably at
30 least 2 to 1 and suitably at most 10 to 1 and preferably at most 1 to 1.

EXAMPLE 1

A strain of *Mortierella alpina* was grown in batch culture in an aqueous medium on a mixture of glucose and yeast autolysate to give a culture containing 54.8 g/l of cells containing 24% w/w of a triglyceride oil. The fatty acid composition of this oil was determined to be 18.6% w/w 5,8,11,14-eicosatetraenoic acid (arachidonic acid).

The cell suspension was dispersed using a high shear mixer (Ultra Turrax, IKA) then fed to a high pressure homogeniser (Rannie Lab homogeniser, APV) operating at a pressure of 400 bar. The resulting cell homogenate was then first mixed with a demulsifying agent (Armogard D5390, Akzo-Nobel, Littleborough) to give a concentration of the demulsifier of 2000 ppm. This aqueous phase was then contacted (at ambient temperature and neutral pH) with hexane by high shear mixing (Ultra Turrax, IKA) to ensure good phase contact. A phase ratio of 3:1 aqueous to solvent was used. The resulting emulsion was separated by centrifugation at 3000 g for 3 minutes. The separated hexane phase was purified from residual fermentation antifoam by passing through a silica gel column (Waters Sep-pak) and evaporated to give the purified triglyceride oil. This oil was analysed for 5,8,11,14-eicosatetraenoic acid content by Gas Chromatography. The overall triglyceride yield from hexane contact of the aqueous phase was 65.7% based on the initial oil content of the microbial cells. The 5,8,11,14-eicosatetraenoic acid content of the oil was 28.1% w/w.

In comparison a sample of fermenter culture was

taken and the cells recovered by filtration. The collected cells were dried by freeze drying then mixed with hexane. 5 g of dried cells were contacted with 25 ml of hexane. The solvent was contacted with the cells for 60 minutes at ambient temperature before the solvent extract was recovered by filtration of the suspension. The solvent was purified of residual fermentation antifoam using a silica gel column (Waters Sep-pak) and evaporated to give a purified triglyceride oil. The overall triglyceride yield was 59% based on the initial oil content of the microbial cells. This was analysed for 5,8,11,14-eicosatetraenoic acid by Gas Chromatography giving a concentration of 7.1% w/w.

This shows that considerable breakdown of the 5,8,11,14-eicosatetraenoic acid had occurred during the extraction of dried cells compared to the extraction from homogenised whole cell culture.

Oil yield was higher from extraction from high pressure homogenised whole cells compared to freeze dried cells and the oil quality was higher in terms of the recovered 5,8,11,14-eicosatetraenoic acid.

EXAMPLE 2

A strain of *Mortierella alpina* was grown in batch culture in an aqueous medium on a mixture of glucose and yeast autolysate to give a culture containing 46.0 g/l of cells containing 51.0% w/w of a triglyceride oil. The fatty acid composition of this oil was determined to be 25.1% w/w 5,8,11,14-eicosatetraenoic acid (arachidonic acid).

The cell suspension was dispersed using a high shear mixer (Ultra Turrax) then fed to a high pressure

homogeniser (Rannie) operating at a pressure of 400 bar. This aqueous phase was then contacted (at ambient temperature and neutral pH) with different quantities of hexane by high shear mixing (Ultra Turrax) to ensure good phase contact. Phase ratios of 1:1, 1:1.5, 1:2, 1:2.5, and 1:3 aqueous to solvent were used. The resulting mixed phases were separated by centrifugation in graduated test tubes at 3000 g for 3 minutes and the recovery of the initial solvent phase measured.

TABLE 1

Phase ratio (aqueous:solvent)	Solvent recovery (%)
3:1	4
2:1	15
1:1	88
1:1.5	95
1:2	97
1:2.5	101
1:3	100

In the case of the sample prepared using a phase ratio of 1:2 aqueous:solvent the solvent phase was recovered and the aqueous phase recontacted with sufficient hexane solvent to give a phase ratio of 1:2 aqueous:solvent. As for the original extraction a high shear mixer was used to ensure good contact of the phases. The pooled hexane phases were evaporated and analysed for total triglyceride content and of 5,8,11,14-eicosatetraenoic acid. The total triglyceride yield was

76% and the 5,8,11,14-eicosatetraenoic acid content of the oil was 29.2% w/w.

EXAMPLE 3

5 A strain of *Thraustochytrium* was grown in batch culture in an aqueous medium on a mixture of glucose, yeast extract, peptone and 20 g/l Na₂SO₄ to give a culture containing 15 g/l of cells containing 30% w/w of a triglyceride oil. The composition of the fatty acids of this oil was determined to be 10% w/w 7, 10, 13, 16, 19-
10 docosahexaenoic acid (DHA).

The cell suspension was passed twice through a high pressure homogenised (Niro-Soavi Panda Laboratory homogeniser) operating at a pressure of 500 bar.

15 This aqueous phase was then contacted (at ambient temperature and neutral pH) with three times the aqueous phase volume of hexane by high shear mixing (Ultra Turrax, IKA) to ensure good phase contact.

The resulting mixed phases separated under gravity and the solvent phase recovered and retained.

20 The aqueous phase was contacted again with hexane (volume equal to three times the aqueous phase volume) and the hexane extract separated by gravity settling.

The pooled hexane phases were evaporated and analysed for total triglyceride content and of 4, 7, 10, 13, 16, 19- docosahexaenoic acid (DHA).
25

The total triglyceride yield was 91% and the 4, 7, 10, 13, 16, 19- docosahexaenoic acid (DHA) content of the oil was 8.7% w/w. The peroxide value (PV) of the oil was measured to be less than 5 milliequivalents of active
30 oxygen per kg.

95TJL05S - MS - 2 July 1996

CLAIMS

- 1 A process in which an oil is extracted from oil containing microorganisms which comprises disintegrating the microorganisms and contacting them in the presence of a water content of at least 70% by weight of that originally present in the cellular material with a water immiscible solvent for the oil, separating the solvent from the microorganisms and recovering the oil from the solvent.
- 2 A process in which an oil is extracted from oil containing microorganisms which comprises disintegrating the microorganisms and contacting them in the presence of substantially all of their water content with a water immiscible solvent for the oil, separating the solvent from the microorganisms and recovering the oil from the solvent.
- 3 A process as claimed in Claim 1 or 2 in which the oil is a triglyceride.
- 4 A process as claimed in any preceding claim in which the oil is oxygen- or heat- sensitive.
- 5 A process as claimed in any preceding claim in which the microorganism is disintegrated in the presence of an aqueous culture medium thereby forming an oil containing medium and contacting the said medium with the solvent.
- 6 A process as claimed in any preceding claim in which the disintegrated organisms are repeatedly contacted with solvent thereby extracting successive quantities of oil therefrom.

- 7 A process as claimed in any preceding claim in which previously unextracted disintegrated microorganisms are extracted with solvent which contains triglyceride extracted from disintegrated microorganisms.
- 8 A process as claimed in Claim 7 which is carried out by counter-current extraction.
- 9 A process as claimed in Claim 8 in which the solvent is less dense than the phase containing the disintegrated microorganism and is fed to a vessel in which extraction occurs at a lower level than the phase containing the disintegrated microorganism.
- 10 A process as claimed in Claim 8 in which the solvent is more dense than the phase containing the disintegrated microorganism and is fed to a vessel in which extraction occurs at a higher level than the phase containing the disintegrated microorganism.
- 11 A process as claimed in Claims 9 or 10 in which the vessel is provided with baffles restricting vertical flow which define contact zones between the baffles and means for stirring material in the contact zones.
- 12 A process as claimed in any preceding claim which comprises separating the solvent by centrifuging.
- 13 A process as claimed in any preceding claim in which the disintegrated microorganisms are contacted with a continuous phase of the solvent.
- 14 A process as claimed in Claim 12 in which during extraction the ratio of solvent phase to other matter present is at least 1 to 1 preferably 1.5 to

1, and more preferably at least 2 to 1 and is suitably at most 10:1 and preferably 5:1.

- 15 A process as claimed in any preceding claim in which the cells are disintegrated by pressure homogenisation.
- 16 A process as claimed in any preceding claim in which the solvent is a hydrocarbon.
- 17 A process as claimed in Claim 15 in which the hydrocarbon is hexane.

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/GB 96/01703

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12P7/64 C11B1/10 C12N1/06 C12N1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12P C11B C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 8413 Derwent Publications Ltd., London, GB; Class D16, AN 84-077725 XP002018398 & JP,A,59 028 480 (NISHIYAMA GUM KK) , 15 February 1984 see abstract	1-3,5,7, 16
A	WO,A,92 13086 (MARTEK CORP) 6 August 1992 see page 11; claims	1
A	PATENT ABSTRACTS OF JAPAN vol. 008, no. 114 (C-225), 26 May 1984 & JP,A,59 028480 (KOUGIYOU KAIHATSU KENKYUSHO:KK;OTHERS: 01), 15 February 1984, see abstract	1

	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 November 1996

Date of mailing of the international search report

22.11.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Delanghe, L

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/GB 96/01703

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GB,A,1 089 093 (THE BRITISH PETROLEUM COMPANY) 1 November 1967 see page 5, line 9 - page 6, line 95 ---	1
A	EP,A,0 522 470 (IDEMITSU PETROCHEMICAL CO) 13 January 1993 see claims -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members:

Internat'l Application No

PCT/GB 96/01703

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9213086	06-08-92	AU-B- 661674	03-08-95
		AU-A- 1235592	27-08-92
		BR-A- 9205519	01-03-94
		CA-A- 2101273	25-07-92
		EP-A- 0568608	10-11-93
		IL-A- 100732	29-06-95
		JP-T- 6505384	23-06-94
		NZ-A- 241358	27-09-94
		OA-A- 9909	15-09-94
GB-A-1089093		BE-A- 653910	02-04-65
		DE-A- 1442058	06-02-69
		NL-A,B 6411478	05-04-65
EP-A-0522470	13-01-93	JP-A- 5308979	22-11-93
		JP-A- 5017796	26-01-93
		AU-B- 640875	02-09-93
		AU-A- 1946792	21-01-93
		DE-D- 69210892	27-06-96
		DE-T- 69210892	02-10-96

Form PCT/ISA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.